

SYMMETRIC-CONVEX FUNCTIONALS OF LINEAR GROWTH

FRANZ GMEINER

ABSTRACT. We discuss existence and regularity theorems for convex functionals of linear growth that depend on the symmetric rather than the full gradients. Due to the failure Korn's Inequality in the L^1 -setup, the full weak gradients of minima do not need to exist, and the paper aims for presenting methods that help to overcome these issues as to partial regularity and higher integrability of minimisers.

1. INTRODUCTION

The purpose of the present paper is to survey and to announce existence and regularity results for minima of autonomous variational integrals which depend on the symmetric rather than the full gradients. More precisely, let Ω be an open and bounded Lipschitz subset of \mathbb{R}^n and consider the variational principle

$$(1.1) \quad \text{to minimise } \mathfrak{F}[v] := \int_{\Omega} f(\varepsilon(v)) \, dx \text{ over a Dirichlet class } \mathcal{D},$$

where $\varepsilon(v) := \frac{1}{2}(Dv + D^T v)$ is the symmetric part of the weak gradient of a function $v: \Omega \rightarrow \mathbb{R}^n$, and $f \in C(\mathbb{R}_{\text{sym}}^{n \times n})$ is a convex function of *linear growth*. By the latter, we understand that there exist $c_1, c_2 > 0$ such that

$$(1.2) \quad c_1 |\mathbf{Z}| \leq f(\mathbf{Z}) \leq c_2(1 + |\mathbf{Z}|) \quad \text{for all } \mathbf{Z} \in \mathbb{R}_{\text{sym}}^{n \times n}.$$

Under these conditions imposed on f , \mathfrak{F} is well-defined on the space $\text{LD}(\Omega)$ consisting of all $v \in L^1(\Omega; \mathbb{R}^n)$ whose weak symmetric gradients belong to $L^1(\Omega; \mathbb{R}_{\text{sym}}^{n \times n})$. This space is equipped with the canonical norm $\|v\|_{\text{LD}} := \|v\|_{L^1} + \|\varepsilon(v)\|_{L^1}$, and we define $\text{LD}_0(\Omega)$ to be the $\|\cdot\|_{\text{LD}}$ -closure of $C_c^1(\Omega; \mathbb{R}^n)$. Consequently, we put $\mathcal{D} := u_0 + \text{LD}_0(\Omega)$ for some fixed $u_0 \in \text{LD}(\Omega)$ and easily conclude that \mathfrak{F} is coercive on \mathcal{D} with respect to the LD-norm.

It is important to note that the aforementioned coerciveness fails when LD is replaced by $W^{1,1}$. The reason for this is the lack of Korn's Inequality, a fundamental obstruction which we briefly describe now. Given $1 < p < \infty$, Korn's Inequality asserts that there exists a constant $C > 0$ such that

$$(1.3) \quad \int_{\Omega} |Dv|^p \, dx \leq C \int_{\Omega} |\varepsilon(v)|^p \, dx$$

holds for all $v \in C_c^1(\Omega; \mathbb{R}^n)$. In consequence, for convex and continuous integrands $g: \mathbb{R}_{\text{sym}}^{n \times n} \rightarrow \mathbb{R}$ which satisfy $c_1 |\mathbf{Z}|^p \leq g(\mathbf{Z}) \leq c_2(1 + |\mathbf{Z}|^p)$ for all $\mathbf{Z} \in \mathbb{R}_{\text{sym}}^{n \times n}$ and two constants $c_1, c_2 > 0$, (1.3) implies that the variational integral $\mathfrak{G}[v] := \int_{\Omega} g(\varepsilon(v)) \, dx$ not only is well-defined on $W^{1,p}(\Omega; \mathbb{R}^n)$ but also coercive on suitable Dirichlet subclasses of $W^{1,p}(\Omega; \mathbb{R}^n)$ with respect

Date: October 25, 2016.

2010 Mathematics Subject Classification. Primary: 49J06; Secondary: 35J06.

Key words and phrases. Functionals of Linear Growth, Regularity Theory, Functions of Bounded Deformation.

to the usual $W^{1,p}$ -norm. As shall be explained in more detail in section 2 below, the reason for (1.3) to hold is that the map $\Phi: \varepsilon(v) \mapsto Dv$ (where $v \in C_c^1(\Omega; \mathbb{R}^n)$) is tacitly identified with its trivial extension to the entire \mathbb{R}^n is a singular integral of convolution type. Thus, by standard results for such operators, Φ is of strong- (p, p) type if and only if $1 < p < \infty$. In turn, if $p = 1$, inequality (1.3) fails to hold true, a fact which is often referred to as *Ornstein's Non-Inequality*. Even stronger statements are available, some of which shall be discussed in section 2.

Since $LD(\Omega)$ is a non-reflexive space, the second chief obstruction is that minimising sequences $(v_k) \subset \mathcal{D}$ might not possess weakly convergent subsequences even though they are uniformly bounded with respect to the LD-norm. To overcome this lack of compactness, it is reasonable to define the space $BD(\Omega)$ of *functions of bounded deformation* as the collection of all $v \in L^1(\Omega; \mathbb{R}^n)$ such that the distributional symmetric gradient $\varepsilon(v)$ can be represented by a $\mathbb{R}_{\text{sym}}^{n \times n}$ -valued Radon measure of finite total variation on Ω , in symbols $\varepsilon(v) \in \mathcal{M}(\Omega; \mathbb{R}_{\text{sym}}^{n \times n})$; see [39, 1] for a detailed treatment of these spaces. In particular, by Ornstein's Non-Inequality, there exist elements $v \in BD(\Omega)$ such that $Dv \notin \mathcal{M}(\Omega; \mathbb{R}^{n \times n})$ and hence $BD(\Omega)$ contains $BV(\Omega; \mathbb{R}^n)$ as a proper subspace. In many respects, the properties of BD-functions are reminiscent of those of BV-functions, and we shall discuss similarities and discrepancies between the two function spaces as the paper evolves. Before passing to criteria that ensure the regularity of minima, we briefly revisit the treatment of the boundary value problem in BD which appears in a similar vein as that in BV as set up in the fundamental work of Giaquinta, Modica and Souček [22].

1.1. Relaxation and Generalised Minima. As an easy consequence of the Banach-Alaoglu and Rellich-Kondrachov Theorems, uniformly bounded sequences in $LD(\Omega)$ possess subsequences that converge to some $v \in BD(\Omega)$ in the weak*-sense. By this we understand that for some $(v_{j(k)}) \subset (v_j)$ there holds $v_{j(k)} \rightarrow v$ strongly in $L^1(\Omega; \mathbb{R}^n)$ and $\varepsilon(v_{j(k)}) \xrightarrow{*} \varepsilon(v)$ in the sense of weak*-convergence of Radon measures on Ω as $k \rightarrow \infty$. In this situation, the weak*-limit map can be shown to exist, however, to establish a reasonable notion of minimality for v , the functional \mathfrak{F} must be extended to $BD(\Omega)$ first. To keep the presentation simple, we stick to the L^1 -Lebesgue-Serrin extension given by

$$\bar{\mathfrak{F}}[v] := \inf \left\{ \liminf_{k \rightarrow \infty} \mathfrak{F}[v_k] : (v_k) \subset \mathcal{D}, v_k \rightarrow v \text{ in } L^1(\Omega; \mathbb{R}^n) \text{ as } k \rightarrow \infty \right\}, \quad v \in BD(\Omega).$$

Note that this type of relaxation is reasonable indeed: If $(v_k) \subset \mathcal{D}$ is bounded with respect to the LD-norm and converges to some $v \in L^1(\Omega; \mathbb{R}^n)$ strongly in $L^1(\Omega; \mathbb{R}^n)$, then $v \in BD(\Omega)$ by lower semicontinuity of the total deformation $|\varepsilon(\cdot)|(\Omega)$ with respect to strong L^1 -convergence. It needs to be noted that the functionals $\bar{\mathfrak{F}}$ admit the explicit integral representation

$$\bar{\mathfrak{F}}[v] = \int_{\Omega} f(\mathcal{E}v) dx + \int_{\Omega} f^{\infty} \left(\frac{dE^s v}{d|E^s v|} \right) d|E^s v| + \int_{\partial\Omega} f^{\infty}(\text{Tr}(v - u_0) \odot \nu_{\partial\Omega}) d\mathcal{H}^{n-1}$$

for $v \in BD(\Omega)$, where

$$\varepsilon(v) = E^{ac}v + E^s v = \frac{dE^c v}{d\mathcal{L}^n} \mathcal{L}^n + \frac{dE^s v}{d|E^s v|} |E^s v| = \mathcal{E}v \mathcal{L}^n + \frac{dE^s v}{d|E^s v|} |E^s v|$$

is the Radon-Nikodým decomposition of $\varepsilon(v)$ into its absolutely continuous and singular parts with respect to Lebesgue measure \mathcal{L}^n ; moreover, $f^{\infty}: \mathbb{R}_{\text{sym}}^{n \times n} \rightarrow \mathbb{R}$ is the *recession*

function of f defined as

$$f^\infty(\mathbf{Z}) := \lim_{t \searrow 0} t f(\mathbf{Z}/t), \quad \mathbf{Z} \in \mathbb{R}_{\text{sym}}^{n \times n},$$

and captures the behaviour of the integrand at infinity, that is to say, where the Lebesgue density of $\varepsilon(v)$ with respect to \mathcal{L}^n becomes singular. Note that by [1, 28], the density $\frac{d\mathbb{E}v}{d\mathcal{L}^n}$ can be shown to equal the symmetric part $\mathcal{E}v$ of the approximate gradient of v \mathcal{L}^n -a.e.. The trace terms in fact are sensible, as by [39, 3], BD-functions attain boundary values in the L^1 -sense. Noting that for $a, b \in \mathbb{R}^n$, $a \odot b := \frac{1}{2}(ab^\top + ba^\top)$ is the symmetric tensor product and $\nu_{\partial\Omega}$ the outer unit normal to $\partial\Omega$, the boundary integral term appearing in the integral representation admits the interpretation of a penalisation term that leads to larger values of the functional provided the L^1 -distance of $\text{Tr}(v)$ from $\text{Tr}(u_0)$ is increased. The proof of the above representation follows along the lines of [7] in the full gradient case, and for more background information, the reader is referred to [26]. For completeness, we make the following

Definition 1 (BD-Minima). An element $u \in \text{BD}(\Omega)$ is called a BD-minimiser if and only if $\mathfrak{F}[u] \leq \mathfrak{F}[v]$ for all $v \in \text{BD}(\Omega)$.

On a sidenote, let us remark that convexity of f substantially simplifies the proof of the integral representation for the relaxed functional. In fact, in the convex setup, it is possible to use the Goffman–Serrin relaxation machinery [27], whereas in the quasiconvex situation more subtle arguments need to be invoked; see the work of Rindler [36] for more detail.

Another notion of minimisers has been employed by Bildhauer & Fuchs [9, 7] in the setting of linear growth functionals on BV, whose adaption to the present situation reads as follows:

Definition 2 (Generalised Minima). Let Ω be an open and bounded Lipschitz subset of \mathbb{R}^n and fix a boundary datum $u_0 \in \text{LD}(\Omega)$. The set of *generalised minima* of \mathfrak{F} given by (1.1) consists of all those $u \in \text{BD}(\Omega)$ for which there exists an \mathfrak{F} -minimising sequence $(u_k) \subset \mathcal{D}_{u_0} := u_0 + \text{LD}_0(\Omega)$ such that $u_k \rightarrow u$ strongly in $L^1(\Omega; \mathbb{R}^n)$ as $k \rightarrow \infty$. The set of all generalised minima is denoted $\text{GM}(\mathfrak{F})$.

Now, if $f \in C(\mathbb{R}_{\text{sym}}^{n \times n})$ is convex – that is to say, f is *symmetric-convex* – then $u \in \text{BD}(\Omega)$ can be shown to be a BD-minimiser if and only if it is a generalised minimiser, and in this case there holds

$$(1.4) \quad \mathfrak{F}[u] = \min \mathfrak{F}(\text{BD}(\Omega)) = \inf \mathfrak{F}[\mathcal{D}].$$

The crucial point thus is to establish existence of a BD-minimiser. This is, however, easily achieved by employing the direct method and by use of Reshetnyak-type theorems on the lower semicontinuity of functionals of measures with respect to the weak*- and strict topologies; see [35, 7]. Hence a satisfactory existence theory is established, and the foremost aim of the present paper is to survey the regularity properties of generalised minima.

1.2. Organisation of the Paper and Description of Results. Having settled existence of generalised minima in the previous section, the paper focusses on regularity results in all of what follows. In section 2, we revisit Korn’s Inequality and Ornstein’s Non-Inequality and strengthen the sketchy arguments outlined above to understand the chief obstructions

for regularity results in the symmetric-convex case. In section 3, we report on recent developments regarding the Hölder and Sobolev regularity of generalised minima of symmetric-convex functionals subject to strong convexity conditions imposed on the variational integrands f . Firstly describing a partial regularity result due to the author [25] in the spirit of Anzellotti & Giaquinta [2], we then turn to conditions on the variational integrands to produce generalised minima of class BV_{loc} or even $W_{\text{loc}}^{1,p}$ for some $1 < p < \infty$, the latter being joint work with Jan Kristensen [26]. To our best knowledge, these results are the first of their kind and extend the regularity theory on the Dirichlet problem on BV to that on BD ; see [7, 5]. To conclude with, in section 4 we introduce the spaces $W^{\mathbb{A},1}$ and $BV_{\mathbb{A}}$ as suitable generalisations of BV and BD and highlight open questions that would lead to a satisfactory existence and regularity theory in this fairly general setup.

ACKNOWLEDGMENT

The author is indebted to the University of Zürich for financial support to attend the 9th European Conference on Elliptic and Parabolic Problems held in Gaeta in May 2016. Moreover, he gratefully acknowledges the comments of an anonymous referee which helped to improve the exposition of the material.

NOTATION

The symmetric $n \times n$ -matrices are denoted $\mathbb{R}_{\text{sym}}^{n \times n}$, and we use the symbol $\langle \cdot, \cdot \rangle$ for the euclidean inner product on finite dimensional spaces. Lastly, \mathcal{L}^n and \mathcal{H}^{n-1} denote the n -dimensional Lebesgue- or $(n-1)$ -dimensional Hausdorff measures, respectively, and we use $(u)_U := \int_U u \, dx := (\mathcal{L}^n(U))^{-1} \int_U u \, dx$ for the mean value of a locally integrable function $u: U \rightarrow \mathbb{R}^N$ whenever this is well-defined.

2. KORN'S INEQUALITY AND ORNSTEIN'S NON-INEQUALITY

Before we embark on the regularity of generalised minima as addressed in the introduction, we briefly wish to comment on Korn's Inequality in slightly more detail. To this end, let $\mathbb{A}[D]$ be a linear, homogeneous first order and constant coefficient differential operator between the two finite-dimensional real vector spaces V and W , i.e., $\mathbb{A}[D]$ can be written in the form

$$(2.1) \quad \mathbb{A}[D] = \sum_{|\alpha|=1} \mathbb{A}_\alpha \partial^\alpha,$$

where $\mathbb{A}_\alpha: V \rightarrow W$ are fixed linear mappings. We associate with $\mathbb{A}[D]$ its *symbol map*

$$(2.2) \quad \mathbb{A}[\xi] := \sum_{|\alpha|=1} \xi_\alpha \mathbb{A}_\alpha, \quad \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n,$$

and call $\mathbb{A}[D]$ *elliptic* if and only if $\mathbb{A}[\xi]: V \rightarrow W$ is injective for any $\xi \neq 0$. Given an elliptic operator $\mathbb{A}[D]$ and $u \in C_c^\infty(\mathbb{R}^n; \mathbb{R}^n)$, we can thus retrieve u from $\mathbb{A}[D]u$ by means of the operator

$$u(x) = \mathbf{G}[\mathbb{A}[D]u](x) = c_n \mathcal{F}_{\xi \mapsto x}^{-1}((\mathbb{A}^*[\xi] \circ \mathbb{A}[\xi])^{-1} \widehat{\mathbb{A}^*[\xi] \mathbb{A}[D]u}) =: \Phi(\mathbb{A}[D]u)(x), \quad x \in \mathbb{R}^n,$$

where $c_n > 0$ is a constant and $\mathbb{A}^*[\xi]$ is the adjoint symbol of $\mathbb{A}[\xi]$ being defined in the obvious manner. Since $(\mathbb{A}^*[\xi] \circ \mathbb{A}[\xi])^{-1} \mathbb{A}^*[\xi]$ is homogeneous of degree -1 , it is easily seen

that \mathbf{G} is a Riesz potential operator of order 1, in particular, we have the bound

$$|\mathbf{G}[v](x)| \lesssim \int_{\mathbb{R}^n} \frac{|v(y)|}{|x-y|^{n-1}} dy \quad \text{for all } x \in \mathbb{R}^n.$$

Finally, differentiating $\mathbf{G}[\mathbb{A}[D]u]$ immediately yields that Du can be written as a singular integral of convolution type acting on $\mathbb{A}[D]u$; see [38]. Therefore, the operator $\Phi: \mathbb{A}[D]u \mapsto Du$ given by $\Phi(\mathbb{A}[D]u) := D(\mathbf{G}[\mathbb{A}[D]u])$ extends to a bounded linear operator $\tilde{\Phi}: L^p(\mathbb{R}^n; W) \rightarrow L^p(\mathbb{R}^n; \mathbb{R}^n \times V)$ provided $1 < p < \infty$. Hence, given $1 < p < \infty$ and an open subset Ω of \mathbb{R}^n , Korn's inequality $\|Du\|_{L^p(\Omega; \mathbb{R}^n \times V)} \leq C\|\mathbb{A}[D]u\|_{L^p(\Omega; W)}$ for all $u \in C_c^\infty(\Omega; V)$ with a finite constant $C = C(\mathbb{A}, p) > 0$ follows from the aforementioned boundedness of singular integrals by extending elements of $C_c^\infty(\Omega; V)$ to \mathbb{R}^n by zero.

Korn's Inequality in the form as given above can be generalised to various other settings; see [32] and [10, 11] for more recent developments in the context of Orlicz functions. Its failure in the case $p = 1$ has witnessed a variety of notable contributions beyond Ornstein's original article [34]; see, among others, [14, 29]. In particular, as can be seen best through Theorem 1.3 of Kirchheim and Kristensen's study [29], there are only trivial L^1 -estimates in the following sense:

Theorem 1 ([29], Theorem 1.3). *Let V, W, X be three finite-dimensional vector spaces and consider two k -th order linear and homogeneous differential operators $\mathbb{A}_1[D]$ and $\mathbb{A}_2[D]$ of the form*

$$\mathbb{A}_1[D] = \sum_{|\alpha|=k} \mathbb{A}_\alpha^1(x) \partial^\alpha \quad \text{and} \quad \mathbb{A}_2[D] = \sum_{|\alpha|=k} \mathbb{A}_\alpha^2(x) \partial^\alpha,$$

with locally integrable coefficients $\mathbb{A}_\alpha^1 \in L_{\text{loc}}^1(\mathbb{R}^n; \mathcal{L}(V; W))$ and $\mathbb{A}_\alpha^2 \in L_{\text{loc}}^1(\mathbb{R}^n; \mathcal{L}(V; X))$ for all $|\alpha| = k$, respectively, the following are equivalent:

- (1) *There exists a constant $C > 0$ such that $\|\mathbb{A}_2[D]\varphi\|_{L^1(\mathbb{R}^n; W)} \leq c\|\mathbb{A}_1[D]\varphi\|_{L^1(\mathbb{R}^n; W)}$ holds for all $\varphi \in C_c^\infty(\mathbb{R}^n; V)$.*
- (2) *There exists $C \in L^\infty(\mathbb{R}^n; \mathcal{L}(W; X))$ with $\|C\|_{L^\infty(\mathbb{R}^n; \mathcal{L}(W; X))} \leq c$ such that $\mathbb{A}_\alpha^2(x) = C(x)\mathbb{A}_\alpha^1(x)$ for \mathcal{L}^n -a.e. $x \in \mathbb{R}^n$ and each $\alpha \in \mathbb{N}_0^n$ with $|\alpha| = k$.*

Here, triviality of L^1 -estimates means that *if* a Korn-type inequality holds (1), then the coefficients are the same up to multiplication with an L^∞ -function. It is important to note that both the symmetric gradient operator $\mathbb{A}_1[D]u = \varepsilon(u)$ or the trace-free symmetric gradient operator $\mathbb{A}_1[D]u = \varepsilon(u) - \frac{1}{n} \operatorname{div}(u) \mathbb{1}_{n \times n}$ with the $(n \times n)$ -unit matrix $\mathbb{1}_{n \times n} \in \mathbb{R}^{n \times n}$ do *not* verify (2) with $\mathbb{A}_2[D] = Du$ and hence they do not admit Korn-type inequalities in the L^1 -setup.

3. PARTIAL AND SOBOLEV REGULARITY

Besides existence of generalised minima as outlined in the introduction, it is natural to investigate their regularity properties, a program which has been launched in the BV-setting in [22, 31]. Here, we focus on Hölder- and Sobolev regularity and shall describe the main obstructions that come along both with the linear growth hypothesis and Ornstein's Non-Inequality. In particular, the latter motivates to study conditions imposed on the variational integrand f that guarantee existence of the full gradients of generalised minima as elements of $\mathcal{M}(\Omega; \mathbb{R}^{n \times n})$ or $L^1(\Omega; \mathbb{R}^{n \times n})$.

3.1. Hölder Regularity. To begin with, let us note that by the genuine vectorial nature of the functional \mathfrak{F} , BD-minima cannot be shown to share everywhere $C^{1,\alpha}$ -regularity unless strong structural conditions are imposed on the variational integrands f . This is in the spirit of the famous counterexamples due to De Giorgi [15] and Giusti & Miranda [24] (see also [33] for an excellent overview) which demonstrate that in the case $N > 1$, functionals of the form

$$\mathfrak{G}[v] := \int_{\Omega} g(\nabla v) \, dx, \quad v: \Omega \rightarrow \mathbb{R}^N$$

do not necessarily produce minimisers of class $C_{\text{loc}}^{1,\alpha}(\Omega; \mathbb{R}^N)$ even if suitable ellipticity, boundedness and measurability assumptions are imposed on the integrands g . The correct substitute is then given by the notion of *partial regularity*: Given $u \in \text{BD}(\Omega)$, we define its *regular set*

$$(3.1) \quad \Omega_u := \{x \in \Omega: \nabla u \text{ is Hölder continuous in a neighbourhood of } x\},$$

and note that the definition of Ω_u depends on the full weak gradients ∇u indeed. We call Ω_u the *regular set* of u and its relative complement $\Sigma_u := \Omega \setminus \Omega_u$ the *singular set*. Adopting these notions, we say that $u \in \text{BD}(\Omega)$ is *partially regular* if and only if Ω_u is open and $\mathcal{L}^n(\Sigma_u) = 0$.

Referring the reader to [4, 23, 33] for a comprehensive overview of techniques to establish partial regularity of minima of elliptic variational integrals, we note that most approaches to the partial regularity rely on the higher integrability of gradients. Such higher integrability results in turn often stem from Caccioppoli-type inequalities in conjunction with Gehring's lemma. Indeed, to sketch the prototype form of such an argument, let $1 < p < 2$ and assume that $u \in W^{1,p}(\Omega; \mathbb{R}^N)$ satisfies a Caccioppoli-type inequality of the form

$$(3.2) \quad \int_{B(z,r)} |Du|^p \, dx \leq C \int_{B(z,2r)} \left| \frac{u - (u)_{B(z,2r)}}{r} \right|^p \, dx$$

for all $z \in \Omega$ and $0 < r < \text{dist}(z, \partial\Omega)/2$. Now, applying the Sobolev–Poincaré-inequality to the right side, we deduce that $\int_{B(z,r)} |Du|^p$ can be locally estimated against $\int_{B(z,2r)} |Du|^q \, dx$ for some $1 < q < p$, and in this sense Du satisfies a reverse Hölder inequality. By Gehring's Lemma, we then obtain that Du belongs to some $L_{\text{loc}}^{p+\varepsilon}$ for some $\varepsilon > 0$, and the reader will notice that the above argument remains unchanged when D is replaced by ε and the mean values on the right side of (3.2) by suitable rigid deformations, that is, elements of the nullspace of ε .

If $p = 1$, then Gehring's Lemma self-improves the integrability of Du or $\varepsilon(u)$, respectively, if and only if we can choose $q < p = 1$. Thus a suitable Sobolev–Poincaré inequality would be required, estimating the L^1 -norm of a function against the L^q -norm of its gradient or symmetric gradient, respectively. However, in the full generality as needed here, this is ruled out by a counterexample due to Buckley & Koskela [13]. In turn, one is lead to the so-called *weak reverse Hölder classes* WRH whose elements satisfy suitable Sobolev–Poincaré inequalities for $q < p = 1$ by definition in a natural way; see [13] for more background information. Coming back to the higher integrability addressed above, it is not clear that the gradients of minima belong to such weak reverse Hölder classes at the relevant stage of the proof; hence different methods are required in the linear growth setting. In the symmetric-convex case as described in the introduction, such estimates can be achieved by use difference quotient-type methods, but in turn require strong ellipticity assumptions on the integrands; see section 3.2 below.

A direct approach to the partial regularity that is particularly designed for convex functionals and applies to functionals of linear growth, too, is that of Anzellotti & Giaquinta [2]. As usual, this particular method also relies on decay estimates for suitable excess functionals too. To describe the decisive feature of this method, let us remark that unlike other, perhaps more standard approaches like blow-up proofs, Anzellotti & Giaquinta derive the required decay estimates through comparing minima with suitable mollifications thereof. These mollifications in turn are shown to be close to solutions of elliptic second order PDE and thus enjoy good decay estimates which then are shown to carry over to the generalised minima themselves. Relying on mollifications and, consequently, Jensen's inequality, the method is well-suited for convex problems, whereas it is not clear how to generalise it to quasiconvex integrands, for instance. With the case of full gradients being treated in [2], the respective generalisation to the symmetric gradient case will be given in [25] by the following

Theorem 2. *Let $f \in C^2(\mathbb{R}_{\text{sym}}^{n \times n}; \mathbb{R}_{\geq 0})$ be convex and of linear growth. Suppose that $u \in \text{BD}(\Omega)$ is a BD-minimiser of \mathfrak{F} given by (1.1). If $(x, z) \in \Omega \times \mathbb{R}_{\text{sym}}^{n \times n}$ is such that*

$$\lim_{R \searrow 0} \left[\int_{B(x, R)} |\mathcal{E}u - z| \, dx + \frac{|E^s u|(B(x, R))}{\mathcal{L}^n(B(x, R))} \right] = 0$$

and $f''(z)$ is positive definite, then $u \in C^{1, \alpha}(U; \mathbb{R}^n)$ for a suitable neighbourhood U of x for all $0 < \alpha < 1$.

Assuming the theorem, the standard Lebesgue differentiation theorem for Radon measures yields the claimed partial regularity: There exists an open subset Ω_u of Ω with $\mathcal{L}^n(\Omega \setminus \Omega_u) = 0$ such that for any $x \in \Omega_u$ there exists $r > 0$ with $u \in C^{1, \alpha}(B(x, r); \mathbb{R}^n)$ for every $0 < \alpha < 1$. We wish to conclude with the following

Remark 1 (Non-Autonomous Integrands). Since \mathfrak{F} given by (1.1) is autonomous, it is possible to overcome the higher integrability of the symmetric gradients in the proof of the above theorem. If the integrand in addition is x -dependent, then the higher integrability seems to be a necessary to conclude the result in this non-autonomous case too. Going back to the discussion at the beginning of the section, such a result is therefore unlikely to be established by means of the method as described above.

3.2. Sobolev Regularity. Next we turn to Sobolev regularity of generalised minima and hereafter aim for conditions on the integrands f under which generalised minima genuinely belong to $\text{BV}_{\text{loc}}(\Omega; \mathbb{R}^n)$ or $W_{\text{loc}}^{1, p}(\Omega; \mathbb{R}^n)$ for some $1 < p < \infty$. To obtain such results, we shall work with a strong convexity adapted from that of Bildhauer & Fuchs [9] in the full gradient case:

Definition 3 (μ -ellipticity). Let $\mu > 1$. A C^2 -integrand $f: \mathbb{R}_{\text{sym}}^{n \times n} \rightarrow \mathbb{R}_{\geq 0}$ is called μ -elliptic if and only if there exist $0 < \lambda \leq \Lambda < \infty$ such that

$$(3.3) \quad \lambda \frac{|\mathbf{A}|^2}{(1 + |\mathbf{B}|^2)^{\frac{\mu}{2}}} \leq \langle f''(\mathbf{B})\mathbf{A}, \mathbf{A} \rangle \leq \Lambda \frac{|\mathbf{A}|^2}{(1 + |\mathbf{B}|^2)^{\frac{1}{2}}}$$

holds for all $\mathbf{A}, \mathbf{B} \in \mathbb{R}_{\text{sym}}^{n \times n}$. We further say that the variational integral \mathfrak{F} is μ -elliptic provided its integrand f is.

Prototypical examples are given by the area-type integrands $m_p(\boldsymbol{\xi}) := (1 + |\boldsymbol{\xi}|^p)^{\frac{1}{p}}$ for $\boldsymbol{\xi} \in \mathbb{R}_{\text{sym}}^{n \times n}$, $1 < p < \infty$; so, for instance, the area integrand m_2 is 3-elliptic, whereas m_p coincides with a $\mu = p + 1$ -elliptic integrand away from the unit ball; also see [9] for more detail. It

is important to remark that $\mu = 1$ is explicitly excluded in definition 3; indeed, 1-elliptic integrands correspond to $L \log L$ -growth. For such integrands, $\varepsilon(u) \in L \log L_{\text{loc}}(\Omega; \mathbb{R}_{\text{sym}}^{n \times n})$ already implies $Du \in L^1_{\text{loc}}(\Omega; \mathbb{R}^{n \times n})$ and hence the full gradients are known to exist and belong to L^1 locally. In this setup of $L \log L$ -growth, the corresponding regularity theory has been established by Fuchs, Seregin and collaborators [17, 19, 18] among others; see the extensive monograph [20] for more information.

3.2.1. Results on the Dirichlet Problem on BV. To explain our method, it is convenient to firstly report on the available higher integrability results for μ -elliptic functionals in the full gradient case

$$(3.4) \quad \mathcal{F}[u] := \int_{\Omega} f(\nabla v) \, dx, \quad v \in \mathcal{D} := u_0 + W_0^{1,1}(\Omega; \mathbb{R}^N),$$

where $u_0 \in W^{1,1}(\Omega; \mathbb{R}^N)$ is a given boundary datum. In analogy with Definition 2, we say that $v \in \text{BV}(\Omega; \mathbb{R}^N)$ is a *generalised minimiser* for \mathcal{F} if and only if there exists an \mathcal{F} -minimising sequence $(v_k) \subset \mathcal{D}$ such that $v_k \rightarrow v$ strongly in $L^1(\Omega; \mathbb{R}^N)$ as $k \rightarrow \infty$.

Using a vanishing viscosity approach, Bildhauer [8] provided the first higher integrability results for gradients of generalised minima. Precisely, assuming $u_0 \in W^{1,2}(\Omega; \mathbb{R}^N)$ for the boundary data, the functional \mathcal{F} is stabilised by adding small Laplacians, i.e., we consider

$$\mathcal{F}_{\delta}[v] := \mathcal{F}[v] + \frac{\delta}{2} \int_{\Omega} |\nabla v|^2 \, dx \quad \text{on } \mathcal{D} := u_0 + W_0^{1,2}(\Omega; \mathbb{R}^N)$$

and finally aim for sending $\delta \searrow 0$. Denoting the unique minimiser of \mathcal{F}_{δ} over \mathcal{D} by u_{δ} , it is easy to prove that (u_{δ}) is a minimising sequence for \mathcal{F} . Bildhauer, in turn building on ideas of Seregin [37], then was able to show that if (u_{δ}) satisfies the *local boundedness assumption*

$$(3.5) \quad \text{for all } K \Subset \Omega \text{ there exists } C(K) > 0 \text{ with } \sup_{0 < \delta < 1} \|u_{\delta}\|_{L^{\infty}(K; \mathbb{R}^N)} \leq C(K),$$

then the weak*-limit u of (u_{δ}) belongs to $W_{\text{loc}}^{1,p}(\Omega; \mathbb{R}^N)$ for some $p = p(\mu) > 1$ provided $1 < \mu < 3$, and to $W_{\text{loc}}^{1,L \log L}(\Omega; \mathbb{R}^N)$ provided $\mu = 3$. Apart from the strong assumptions made on the particular minimising sequence, the boundary data and the functional itself, the result merely applies to one particular such generalised minimiser. The reason for this is the possible non-uniqueness of generalised minima which, in turn, is a consequence of the recession parts in the relaxed variational integral. Indeed, even if $f \in C^2(\mathbb{R}^{N \times n})$ is strictly convex, the recession function $f^{\infty}: \mathbb{R}^{N \times n} \rightarrow \mathbb{R}$ is positively 1-homogeneous and thus never strictly convex. In consequence, if a minimiser does not have vanishing singular part with respect to Lebesgue measure, uniqueness in general fails as f and f^{∞} act on two mutually singular parts of the gradients. To achieve uniqueness, one must therefore genuinely rule out the singular parts of minima. This has been achieved recently by Beck & Schmidt [5] by sophisticated use of the Ekeland variational principle in the negative Sobolev space $W^{-1,1}$ (see Prop. 1 below) for the borderline case $\mu = 3$. Referring the reader for the precise outline to [5], the general streamline is this: Starting from an arbitrary minimising sequence $(u_k) \subset u_0 + W_0^{1,1}(\Omega; \mathbb{R}^N)$, the Ekeland variational principle yields another minimising sequence (v_k) that is $W^{-1,1}$ -close to (u_k) and has the same weak*-limit. This new sequence (v_k) is then shown to be a sequence of almost minimisers to suitably stabilised functionals and thus can be proved to belong to $W_{\text{loc}}^{2,2}$. At this point it is possible to adapt Bildhauer's approach and hence, by arbitrariness of (u_k) , uniqueness and the aforementioned higher regularity results follow at once. However, it needs to be stressed

that Beck & Schmidt's method of proof also relies on a version of the local boundedness assumption. Further, assuming Uhlenbeck, i.e., radial structure of the integrands, stronger results such as $C^{1,\alpha}$ -regularity of generalised minima of the \mathcal{F} can be achieved; see [7, 8, 6].

3.2.2. Results on the Dirichlet Problem on BD. Going back to functionals \mathfrak{F} as given by (1.1), the main difficulty lies in the appearance of the *full difference quotients* when aiming for higher integrability estimates and testing the Euler–Lagrange equation of a suitably stabilised functional with the canonical choice $\varphi := \Delta_{s,h}^-(\rho^2 \Delta_{s,h}^+ v)$, where $\Delta_{s,h}^\pm v(x) := \frac{1}{h}(v(x \pm h e_s) - v(x))$ are the forward or backward difference quotients, respectively. By Ornstein's Non-Inequality, $\Delta_{s,h}^+ v$ cannot even be bounded locally in L^1 for $v \in \text{BD}$ in general, and thus the suitable device hence is to work with finite differences instead of difference quotients and to establish estimates for carefully chosen Besov-norms of the symmetric gradients.

We pass on to a more precise description of the method which is, to some extent, inspired by [5]. Let $(v_k) \subset \mathcal{D}$ be a minimising sequence for the μ -elliptic functional \mathfrak{F} given by (1.1), with μ to be determined later on. Then we consider for a suitable sequence $(\alpha_k) \subset \mathbb{R}_{>0}$ with $\alpha_k \searrow 0$ as $k \rightarrow \infty$ the stabilised functionals

$$\mathfrak{F}_k[w] := \int_{\Omega} f(\varepsilon(w)) \, dx + \alpha_k \int_{\Omega} (1 + |\varepsilon(w)|^p) \, dx =: \int_{\Omega} f_k(\varepsilon(w)) \, dx$$

with $p \geq n$ on appropriately modified Dirichlet classes \mathcal{D}_k to keep track of the fact that the leading part of \mathfrak{F}_k is the p -th Dirichlet energy; indeed, as $p > 1$, minima of \mathfrak{F}_k belong to $W^{1,p}$ and thus possess full gradients in L^p by Korn's Inequality. Extending each \mathfrak{F}_k to $(W_0^{1,\infty}(\Omega; \mathbb{R}^n))^*$ by infinity on $(W_0^{1,\infty}(\Omega; \mathbb{R}^n))^* \setminus \mathcal{D}_k$, we obtain a lower semicontinuous functional on $(W_0^{1,\infty}(\Omega; \mathbb{R}^n))^*$ which is continuous with respect to the norm topology on $(W_0^{1,\infty}(\Omega; \mathbb{R}^n))^*$. To continue, we recall the following instrumental

Proposition 1 (Ekeland's Variational Principle). *Let (X, d) be a complete metric space and assume that $f: X \rightarrow [0, \infty]$ is lower semicontinuous with $\inf F(X) < \infty$. If for some $\varepsilon > 0$ and $x \in X$ there holds $F[x] \leq \inf F(X) + \varepsilon$, then there exists $v \in X$ such that $d(x, v) \leq \sqrt{\varepsilon}$ and*

$$F[v] \leq F[w] + \sqrt{\varepsilon} d(v, w) \quad \text{for all } w \in X.$$

For a proof and a discussion of this result, see [23], Thm. 5.6.. Using suitable approximations and Proposition 1 in the Banach space $(W_0^{1,\infty})^*$, we obtain a sequence (u_k) , each of whose members is an almost minimiser of \mathfrak{F} , is $(W_0^{1,\infty})^*$ -close to v_k and, most crucially, each u_k satisfies the perturbed Euler–Lagrange equation

$$\left| \int_{\Omega} \langle f'_k(\varepsilon(u_k)), \varepsilon(\varphi) \rangle \, dx \right| \leq \frac{1}{k} \|\varphi\|_{(W_0^{1,\infty}(\Omega; \mathbb{R}^n))^*} + (\text{small perturbation})$$

for all $\varphi \in W_0^{1,p}(\Omega; \mathbb{R}^n)$. Given $x_0 \in \Omega$, $0 < r < R < \text{dist}(x_0, \partial\Omega)$, we consider for an arbitrary standard unit vector e_s , $s = 1, \dots, n$, the test functions $\varphi := \tau_{s,h}^-(\rho^2 \tau_{s,h}^+ u_k)$ with $\tau_{s,h}^\pm = h \Delta_{s,h}^\pm$. Essentially following, e.g., [23], section 10.1, and employing μ -ellipticity of f , we end up with a coercive inequality

$$(3.6) \quad \int_{\Omega} \frac{|\rho \tau_{s,h}^+ \varepsilon(u_k)|^2}{(1 + |\varepsilon(u_k)|)^\mu} \, dx \lesssim \left| \int_{\Omega} \langle f'_k(\varepsilon(u_k)), \rho \odot \tau_{s,h}^+ u_k \rangle \, dx \right| + \frac{1}{k} \|\tau_{s,h}^-(\rho^2 \tau_{s,h}^+ u_k)\|_{(W_0^{1,\infty})^*},$$

with the constants implicit in ' \lesssim ' being uniformly bounded in k . Let us briefly explain how to handle the two terms on the right side: As to the first term, we note that $f'_k(\varepsilon(u_k))$ converges in a suitable sense to the solution of the dual problem associated with (1.1) (in the sense of convex duality, see [16]). Since the dual solution $\sigma \in L^\infty(\Omega; \mathbb{R}^{n \times n})$ itself belongs to $W_{\text{loc}}^{1,2}(\Omega; \mathbb{R}^{n \times n})$, this regularity can be shown to inherit to $f'_k(\varepsilon(u_k))$ uniformly in k . If we wish to fruitfully use this estimate, we need to suitably bound $\|\rho\tau_{s,h}^+ u_k\|_{L^2}$ uniformly in k too. For general n , the fractional Sobolev-type embeddings $\text{BD}_{\text{loc}} \hookrightarrow W_{\text{loc}}^{s,n/(n-1+s)}$, $0 < s < 1$, with the fractional Sobolev spaces $W^{\theta,r}$, $0 < \theta < 1$ and $r \geq 1$, are optimal, and $n/(n-1+s) = 2$ is achieved if and only if $n = 2$ and $s = 0$. In this case, however, we lose all smoothness information and hence may invoke a condition that is slightly weaker than a local boundedness assumption in the spirit of (3.5); namely, we require the so-called *local BMO-assumption*, meaning that for each relatively compact $K \subset \Omega$, the seminorms $\|u_k\|_{\text{BMO}(K; \mathbb{R}^n)}$ are bounded independently of k . As shall be demonstrated in [26] by means of so-called Dorronsoro-type estimates which have been fruitfully used in [30] by Kristensen & Mingione in a different context, we have

$$(3.7) \quad \text{BD}_{\text{loc}} \cap \text{BMO}_{\text{loc}} \hookrightarrow W_{\text{loc}}^{\frac{1}{p}-\varepsilon,p}$$

for all $1 \leq p < \infty$ and suitably small $\varepsilon > 0$; the limiting case $\varepsilon = 0$ is not even reached in general even if BD_{loc} is replaced by the considerably smaller space $W_{\text{loc}}^{1,1}$, a fact which has been pointed out by Bourgain, Brezis & Mironescu in [12], Remark 3. The upshot of this in comparison with the aforementioned embedding without the BMO-side constraint is that although the BMO-condition is not reflected by the first derivatives, it improves both fractional differentiability and the corresponding integrability at a uniform rate. Putting $p = 2$ in (3.7), it is possible to estimate the first term on the right side of (3.6) by $Ch^{3/2-\varepsilon}$ for any suitably small $\varepsilon > 0$. The second term on the right side of (3.6) can be estimated in the same way, using $\|\Delta_{s,h}^+ v\|_{(W_0^{1,\infty})^*} \lesssim \|v\|_{L^1}$ together with $\text{BD}_{\text{loc}} \hookrightarrow W_{\text{loc}}^{\theta,1}$ for any $0 < \theta < 1$. Going back to (3.6), we then obtain the uniform bound

$$\int_{\Omega} \left| \frac{\rho\tau_{s,h}^+ \varepsilon(u_k)}{h^{\frac{3}{4}-\frac{\varepsilon}{2}}} \right|^2 \omega_k \, dx := \int_{\Omega} \left| \frac{\rho\tau_{s,h}^+ u_k}{h^{\frac{3}{4}-\frac{\varepsilon}{2}}} \right|^2 \frac{1}{(1 + |\varepsilon(u_k)|)^\mu} \, dx \leq C$$

This is a weighted Nikolskiĭ-type estimate for $\varepsilon(u_k)$. At this stage of the proof, it is possible to deduce that the weights ω_k uniformly belong to certain Muckenhoupt classes A_p for some $1 < p < \infty$ and hence, using embedding results for weighted Nikolskiĭ spaces and the theory of singular integrals on Muckenhoupt weighted Lebesgue spaces, it is possible to deduce for a non-empty range of $\mu \in (1, 2)$ that the symmetric gradients $\varepsilon(u_k)$ are uniformly bounded in certain non-weighted L_{loc}^p -spaces with $p > 1$. By arbitrariness of the initially chosen minimising sequence, this implies $u \in W_{\text{loc}}^{1,p}(\Omega; \mathbb{R}^n)$ by Korn's Inequality and thus establishes the higher integrability of generalised minima of (1.1) subject to the above local BMO-assumption. In summary, the strategy leads to the following theorem which shall be established in [26]:

Theorem 3. *Let $f \in C^2(\mathbb{R}_{\text{sym}}^{n \times n})$ be a convex integrand of linear growth and Ω an open and bounded Lipschitz subset of \mathbb{R}^n . For every $n \in \mathbb{N}$ with $n \geq 2$ there exists a number $1 < \mu(n) < 2$ and an exponent $p \geq 1$ such that the following holds: If f is μ -elliptic with $\mu \leq \mu(n)$, then any generalised minimiser $u \in \text{BD}(\Omega) \cap \text{BMO}_{\text{loc}}(\Omega; \mathbb{R}^n)$ of the functional \mathfrak{F} given by (1.1) belongs to $W_{\text{loc}}^{1,p}(\Omega; \mathbb{R}^n)$ for some $p \geq 1$.*

Let us remark that it is plausible for the preceding result to hold true for a fairly larger range of μ than described in Theorem 3, however, this seems hard to be achieved by use of the above method. In particular, a Sobolev regularity result regarding the limiting case $\mu = 3$ (an instance of which is the area integrand $f(\mathbf{Z}) := \sqrt{1 + |\mathbf{Z}|^2}$) would be desirable. Finally, the strategy outlined above allows to weaken even the local BMO–assumption in view of uniform local L^p –bounds on the single members of minimising sequences, and shall be addressed in a future publication.

4. \mathbb{A} –CONVEX FUNCTIONALS

It is a natural extension of the problems and results outlined in the previous sections to replace the symmetric gradient operator by an elliptic differential operator of the form (2.1). In turn, within the framework of section 2, we aim for existence and regularity properties for minima functionals of the form

$$\mathfrak{F}[v] := \int_{\Omega} f(\mathbb{A}[D]u) \, dx$$

over Dirichlet classes $\tilde{\mathcal{D}} := u_0 + W_0^{\mathbb{A},1}(\Omega)$, where $f: W \rightarrow \mathbb{R}$ is of linear growth, thus verifying (1.2) with the obvious modifications. The spaces $W^{\mathbb{A},1}(\Omega)$ are defined similarly to $W^{1,1}$ or LD, namely, we say that a measurable map $v: \Omega \rightarrow V$ belongs to $W^{\mathbb{A},1}(\Omega)$ if and only if $v \in L^1(\Omega; V)$ and the weak differential expression $\mathbb{A}[D]u$ belongs to $L^1(\Omega; W)$. Equipped with the canonical norm, one defines $W_0^{\mathbb{A},1}(\Omega)$ as the closure of $C_c^1(\Omega; V)$ with respect to the norm topology. Similarly, we define the space of functions of bounded $\mathbb{A}[D]$ –variation $BV_{\mathbb{A}}(\Omega)$ to consist of all $v \in L^1(\Omega; V)$ for which the distributional differential expression $\mathbb{A}[D]v$ can be represented by a finite W –valued Radon measure. Based on the linear growth assumption, the direct method in conjunction with the obvious changes of section 1.1 would lead to a satisfactory existence theory within the class of functions bounded $\mathbb{A}[D]$ –variation *provided* the trace operator for such function spaces would be well–understood. This, however, is not the case by now: For instance, even if $\mathbb{A}[D]$ is an elliptic operator in the sense of section 2, elements of $BV_{\mathbb{A}}$ or $W^{\mathbb{A},1}$ do not necessarily possess traces in $L^1(\partial\Omega; V)$ for arbitrary Lipschitz domains $\Omega \subset \mathbb{R}^n$:

Remark 2. For $n = 2$, let the trace–free symmetric gradient operator defined by $\varepsilon^D(v) := \varepsilon(v) - \frac{1}{2} \operatorname{div}(v) \mathbb{1}_{2 \times 2}$ with the (2×2) –unit matrix. In this situation, ε^D is an elliptic, linear, homogeneous first order differential operator on \mathbb{R}^2 from $V = \mathbb{R}^2$ to $W = \mathbb{R}^{2 \times 2}$. Identifying $\mathbb{R}^2 \cong \mathbb{C}$, it is easy to see that $\ker(\varepsilon^D)$ contains all holomorphic functions. To argue that elements of $W^{\varepsilon^D,1}(B(0,1); \mathbb{R}^2) (\simeq W^{\varepsilon^D,1}(\mathbb{D}; \mathbb{C}))$ with the unit disk $\mathbb{D} \subset \mathbb{C}$ do not have traces in $L^1(\partial B(0,1); \mathbb{R}^2) (\simeq L^1(\partial\mathbb{D}; \mathbb{C}))$, consider $f: \mathbb{D} \ni z \mapsto 1/(z-1) \in \mathbb{C}$. Then $f \in L^1(\mathbb{D}; \mathbb{C})$, is holomorphic and thus belongs to $W^{\varepsilon^D,1}(\mathbb{D}; \mathbb{C})$ whereas it is easy to see that $\int_{\partial\mathbb{D}} |f(z)| \, dz = \infty$.

The previous example is due to Fuchs & Repin [21], and motivates the characterisation of all $\mathbb{A}[D]$ such that the corresponding spaces $W^{\mathbb{A},1}(\mathbb{D}; \mathbb{C})$ possess trace space $L^1(\partial\Omega; V)$ at least for the large class of bounded Lipschitz subsets Ω of \mathbb{R}^n . For such operators, the Sobolev regularity result, Theorem 3 is easily shown to hold true as its proof does not use the specific structure of the symmetric gradient operator. In this respect, it is important to note that the techniques available in the literature – so for instance Strang & Temam’s approach [39] in the case of BD – which are tailored for the symmetric gradient case, do not apply to

the setting of arbitrary elliptic differential operators of the form (2.1) without substantial modifications. We hope to successfully tackle this problem in a future publication.

REFERENCES

- [1] Ambrosio, L.; Coscia, A.; Dal Maso, G.: Fine properties of functions with bounded deformation. *Arch. Rational Mech. Anal.* 139 (1997), no. 3, 201–238.
- [2] Anzellotti, G.; Giaquinta, M. Convex functionals and partial regularity. *Arch. Rational Mech. Anal.* 102 (1988), no. 3, 243–272.
- [3] Babadjian, J–F.: Traces of functions of bounded deformation. *Indiana Univ. Math. J.* 64 (2015), no. 4, 1271–1290.
- [4] Beck, L.: Elliptic regularity theory. A first course. *Lecture Notes of the Unione Matematica Italiana*, 19. Springer, Cham; Unione Matematica Italiana, Bologna, 2016. xii+201 pp.
- [5] Beck, L.; Schmidt, T.: On the Dirichlet problem for variational integrals in BV. *J. Reine Angew. Math.* 674 (2013), 113–194.
- [6] Beck, L.; Schmidt, T.: Interior gradient regularity for BV minimizers of singular variational problems. *Nonlinear Anal.* 120 (2015), 86–106.
- [7] Bildhauer, M.: Convex variational problems. Linear, nearly linear and anisotropic growth conditions. *Lecture Notes in Mathematics*, 1818. Springer-Verlag, Berlin, 2003. x+217 pp.
- [8] Bildhauer, M.: A priori gradient estimates for bounded generalised solutions of a class of variational problems with linear growth. *J. Convex Ana.* 9 (2002), 117–137.
- [9] Bildhauer, M.; Fuchs, M.: On a class of variational integrals with linear growth satisfying the condition of μ -ellipticity. *Rend. Mat. Appl.* (7) 22 (2002), 249–274 (2003).
- [10] Breit, D.; Diening, L.: Sharp conditions for Korn inequalities in Orlicz spaces. *J. Math. Fluid Mech.* 14 (2012), no. 3, 565–573.
- [11] Breit, D.; Cianchi, A.; Diening, L.: Trace-free Korn inequalities in Orlicz spaces. Preprint. arXiv:1605.01006
- [12] Bourgain, J.; Brezis, H.; Mironescu, P.: Lifting in Sobolev spaces. *J. Anal. Math.* 80 (2000), 37–86.
- [13] Buckley, S.M.; Koskela, P.: Sobolev-Poincaré inequalities for $p < 1$. *Indiana Univ. Math. J.* 43 (1994), no. 1, 221–240.
- [14] S. Conti; D. Faraco; F. Maggi; S. Müller: Rank-one convex functions on 2×2 - symmetric matrices and laminates on rank-three lines. *Calc. Var. Partial Differential Equations* 24 (2005), no. 4, 479–493.
- [15] De Giorgi, E.: Un esempio di estremali discontinue per un problema variazionale di tipo ellittico. (Italian) *Boll. Un. Mat. Ital.* (4) 1 1968, 135–137.
- [16] Ekeland, I.; Temam, R.: Convex analysis and variational problems. North-Holland, Amsterdam 1976.
- [17] Frehse, J.; Seregin, G.: Regularity of solutions to variational problems of the deformation theory of plasticity with logarithmic hardening. *Proc. St. Petersburg Math. Soc.* 5 (1998), 184–222. English Translation: *Amer. Math. Soc. Transl. II*, 193 (1999), 127–152.
- [18] Fuchs, M.; Mingione, G.: Full $C^{1,\alpha}$ -regularity for free and constrained local minimizers of elliptic variational integrals with nearly linear growth. *Manus. Math.* 102 (2000), 227–250.
- [19] Fuchs, M.; Seregin, G.: A regularity theory for variational integrals with $L\log L$ -growth. *Calc. Var.* 6 (1998), 171–187.
- [20] Fuchs, M.; Seregin, G.: Variational methods for problems from plasticity theory and for generalised Newtonian fluids. *Lecture Notes in Mathematics* 1749, Springer, Berlin–Heidelberg 2000.
- [21] Fuchs, M.; Repin, S. Some Poincaré-type inequalities for functions of bounded deformation involving the deviatoric part of the symmetric gradient. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 385 (2010), Kraevye Zadachi Matematicheskoi Fiziki i Smezhnye Voprosy Teorii Funktsii. 41, 224–233, 237; reprinted in *J. Math. Sci. (N.Y.)* 178 (2011), no. 3, 367–372.
- [22] Giaquinta, M.; Modica, G.; Souček, J.: Functionals with linear growth in the calculus of variations. I, II. *Comment. Math. Univ. Carolin.* 20 (1979), no. 1, 143156, 157–172.
- [23] Giusti, E.: Direct methods in the calculus of variations. World Scientific Publishing Co., Inc., River Edge, NJ, 2003. viii+403 pp.
- [24] Giusti, E.; Miranda, M.: Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni. (Italian) *Boll. Un. Mat. Ital.* (4) 1 1968 219–226.
- [25] Gmeineder, F.: Partial Regularity for Symmetric-Convex Functionals. In Preparation.
- [26] Gmeineder, F.; Kristensen, J.: Sobolev Regularity for Symmetric-Convex Functionals. In Preparation.

- [27] Goffman, C.; Serrin, J.: Sublinear functions of measures and variational integrals. *Duke Math. J.* 31 (1964), 159–178.
- [28] Hajlasz, P.: On approximate differentiability of functions with bounded deformation. *Manuscripta Math.* 91 (1996), no. 1, 61–72.
- [29] Kirchheim, B.; Kristensen, J.: On Rank One Convex Functions that are Homogeneous of Degree One. *Arch. Ration. Mech. Anal.* 221 (2016), no. 1, 527–558.
- [30] Kristensen, J.; Mingione, G.: The singular set of Lipschitzian minima of multiple integrals. *Arch. Ration. Mech. Anal.* 184 (2007), no. 2, 341–369.
- [31] Ladyzhenskaya, O.A.; Ural'tseva, N.N.: Local estimates for gradients of solutions of non-uniformly elliptic and parabolic problems. *Comm. Pure Appl. Math.* 23 (1970), 667–703.
- [32] Málek, J.; Nečas, J.; Rokyta, M.; Ružička, M.: Weak and measure-valued solutions to evolutionary PDEs. *Applied Mathematics and Mathematical Computation*, 13. Chapman & Hall, London, 1996. xii+317 pp.
- [33] Mingione, G.: Regularity of minima: an invitation to the dark side of the calculus of variations. *Appl. Math.* 51 (2006), no. 4, 355–426.
- [34] Ornstein, D.: A non-equality for differential operators in the L^1 -norm. *Arch. Rational Mech. Anal.* 11 1962 40–49.
- [35] Reshetnyak, Yu. G.: Weak convergence of completely additive vector functions on a set. *Siberian Mathematical Journal* (1968), Volume 9, Issue 6, 1039–1045.
- [36] Rindler, F.: Lower Semicontinuity for integral functionals in the space of functions of bounded deformation via rigidity and Young measures. *Arch. Ration. Mech. Anal.* 202 (2011), no. 1, 63–113.
- [37] Seregin, G.: On differential properties of extremals of variational problems arising in plasticity theory. *Differentsial'nye Uravneniya* 26 (1990), 1033–1043 (in Russian). English translation: *Differential Equations* 26 (1990), 756–766.
- [38] Stein, E.M.: Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals. With the assistance of Timothy S. Murphy. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, 1993.
- [39] Strang, G.; Temam, R.: Functions of bounded deformation. *Arch. Rational Mech. Anal.* 75 (1980/81), no. 1, 7–21.

MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, OX2 6HG OXFORD, UNITED KINGDOM

E-mail address: gmeineder@maths.ox.ac.uk

URL: <https://www.maths.ox.ac.uk/people/franz.gmeineder>